

THE JOURNAL OF BONE & JOINT SURGERY

JB&JS

This is an enhanced PDF from The Journal of Bone and Joint Surgery

The PDF of the article you requested follows this cover page.

Early Mortality and Cardiorespiratory Failure in Patients with Fibrodysplasia Ossificans Progressiva

Frederick S. Kaplan, Michael A. Zasloff, Joseph A. Kitterman, Eileen M. Shore, Charles C. Hong and David M. Roche

J Bone Joint Surg Am. 2010;92:686-691. doi:10.2106/JBJS.I.00705

This information is current as of September 28, 2010

Supporting data

<http://www.ejbjs.org/cgi/content/full/92/3/686/DC1>

Reprints and Permissions

Click here to [order reprints or request permission](#) to use material from this article, or locate the article citation on jbjs.org and click on the [Reprints and Permissions] link.

Publisher Information

The Journal of Bone and Joint Surgery
20 Pickering Street, Needham, MA 02492-3157
www.jbjs.org

Early Mortality and Cardiorespiratory Failure in Patients with Fibrodysplasia Ossificans Progressiva

By Frederick S. Kaplan, MD, Michael A. Zasloff, MD, PhD, Joseph A. Kitterman, MD, Eileen M. Shore, PhD, Charles C. Hong, MD, PhD, and David M. Rocke, PhD

Investigation performed at the University of Pennsylvania School of Medicine, Philadelphia, Pennsylvania

Background: Fibrodysplasia ossificans progressiva, a rare genetic disorder of progressive extraskeletal ossification, is the most disabling form of heterotopic ossification in humans. However, little is known about the lifespan or causes of mortality in these patients. We undertook this study to determine the lifespan and causes of mortality in individuals who had fibrodysplasia ossificans progressiva.

Methods: We reviewed comprehensive mortality reports from two large registries of patients with fibrodysplasia ossificans progressiva. Together, these registries comprise >90% of all known patients with this condition in the world. We noted the sex, dates of birth and death, and the cause of death for each individual. We verified the cause of death with extensive medical records, when available. We also collected date of birth, current age, and sex information for each living patient member of the International Fibrodysplasia Ossificans Progressiva Association.

Results: Sixty deaths (thirty male and thirty female patients) were reported in the fibrodysplasia ossificans progressiva community during a thirty-three-year-period. For all sixty patients, the median age at the time of death was forty years (range, three to seventy-seven years). Data were sufficient to establish the cause of death in forty-eight (80%) of the sixty individuals. The median age at the time of death for the forty-eight patients (twenty-four male and twenty-four female patients) with an established cause of death was also forty years. The median lifespan estimated from the 371 individuals in the international fibrodysplasia ossificans progressiva community who were alive and the sixty who had died was fifty-six years (95% confidence interval, fifty-one to sixty years). The most common causes of death in patients with fibrodysplasia ossificans progressiva were cardiorespiratory failure from thoracic insufficiency syndrome (54%; median age, forty-two years) and pneumonia (15%; median age, forty years).

Conclusions: Fibrodysplasia ossificans progressiva is not only an extremely disabling disease but also a condition of considerably shortened lifespan. The most common cause of death in patients with fibrodysplasia ossificans progressiva is cardiorespiratory failure from thoracic insufficiency syndrome.

Fibrodysplasia ossificans progressiva (Mendelian Inheritance in Man [MIM] #135100) is a rare, disabling genetic disorder of progressive heterotopic ossification caused by a heterozygous activating mutation of the gene encoding activin receptor A type I/activin-like kinase 2 (ACVR1/ALK2), a bone morphogenetic protein (BMP) type-I receptor, in all classically affected individuals worldwide¹⁻⁴. There is no racial, ethnic, sex, or geographic predilection⁵.

While disease flare-ups in fibrodysplasia ossificans progressiva are episodic, disability is cumulative^{3,6}. Most patients are confined to a wheelchair by the third decade of life^{6,7}. Clinical

management is symptomatic, and there is no effective treatment^{3,8}. While much is known about the disabling complications of this disease, documentation of lifespan and causes of death in these patients have not been examined. We undertook a comprehensive study to determine the lifespan and cause of death in individuals with fibrodysplasia ossificans progressiva.

Materials and Methods

Mortality Records

We reviewed the comprehensive mortality records from the International Fibrodysplasia Ossificans Progressiva

Disclosure: In support of their research for or preparation of this work, one or more of the authors received, in any one year, outside funding or grants in excess of \$10,000 from the Center for Research in Fibrodysplasia Ossificans Progressiva and Related Disorders, the Ian Cali Endowment, the Weldon Family Endowment, the Isaac and Rose Nassau Professorship of Orthopaedic Molecular Medicine, the International Fibrodysplasia Ossificans Progressiva Association, the Rita Allen Foundation, and the National Institutes of Health (R01-AR40196 and R01-HG003352). Neither they nor a member of their immediate families received payments or other benefits or a commitment or agreement to provide such benefits from a commercial entity.




Fig. 1
Age distribution (in years) at the time of death for sixty patients with fibrodysplasia ossificans progressiva (1973 to 2006).

Association from its inception in 1988 through 2006, and from the International Fibrodysplasia Ossificans Progressiva Clinic at the University of Pennsylvania from its inception in 1973 through 2006. These two patient registries comprise >90% of the known patients with fibrodysplasia ossificans progressiva in the world. All patients were well known to at least one of the investigators and had a confirmed history of fibrodysplasia ossificans progressiva based on the presence of the two classic clinical criteria: congenital malformations of the great toes and progressive heterotopic ossification in characteristic anatomic patterns. Medical records were reviewed to establish the dates of birth and death and the cause of death for each individual. The study was granted an exemption by the institutional review board of the University of Pennsylvania.

Living Individuals

The membership records of the International Fibrodysplasia Ossificans Progressiva Association were reviewed for 2006, in order to provide a concurrent cross-sectional profile of living patients with the condition. The birth year and sex were noted for each individual with fibrodysplasia ossificans progressiva who was living in January 2006.

Statistical Analysis

Kaplan-Meier survival curves were constructed for the merged population of living and deceased individuals with fibrodysplasia ossificans progressiva with use of the R survival package^{9,10}. The survfit function was used to construct the curve and delineate 95% confidence intervals^{9,10}. A comparison survival curve was calculated from mortality data from the U.S. Census (1980) contained in the survexp.us¹⁰. The 1980 U.S. Census was prospectively chosen on the basis of two criteria: the need for: (1) a large well-documented control population within the study period, and (2) a multiethnic control population that best represented the geographic, ethnic, and racial diversity of the study population. Significant differences were analyzed by the log-rank test or by analysis of variance.

Source of Funding

This work was funded in part by the Center for Research in Fibrodysplasia Ossificans Progressiva and Related Disorders, the Ian Cali Endowment, the Weldon Family Endowment, the Isaac and Rose Nassau Professorship of Orthopaedic Molecular Medicine, and by grants from the International Fibrodysplasia Ossificans Progressiva Association, the Rita Allen Foundation, and the National Institutes of Health (R01-AR40196 and R01-HG003352).

Results

There were sixty reported deaths (thirty male and thirty female patients) in the fibrodysplasia ossificans progressiva study population during the thirty-three-year period (1973 to 2006). Data were sufficient to establish an unequivocal cause of death in forty-eight (80%) of sixty individuals (see Appendix). The median age at the time of death for these forty-eight patients (twenty-four male and twenty-four female patients) was forty years (range, three to sixty years), the same median age (forty years; range, three to seventy-seven years) at the time of death for all sixty patients (Fig. 1).

The Kaplan-Meier survival curve estimate from the combined data set of 371 living and sixty deceased individuals with fibrodysplasia ossificans progressiva revealed an estimated median life expectation of fifty-six years (95% confidence interval, fifty-one to sixty years) (Fig. 2). Substantial excess mortality in patients with fibrodysplasia ossificans progressiva began at about the age of thirty years, with only an estimated 30% of the patients with fibrodysplasia ossificans progressiva surviving to the age of sixty years. In contrast, approximately




Fig. 2
Kaplan-Meier survival curves for patients with fibrodysplasia ossificans progressiva (FOP) and for the 1980 U.S. population. The dotted lines are 95% confidence limits. The solid red line represents the actual survival data (average of male and female) for the U.S. population from the U.S. Census Bureau (1980). The Y axis is the percent survival $\times 10^{-2}$, and the X axis is the age in years.

TABLE I Causes of Mortality in Patients with Fibrodysplasia Ossificans Progressiva

Cause	No.	Percentage	M	F	Age Range (yr)	Median Lifespan (yr)
Cardiorespiratory failure	26	54	12	14	8-58	42
Pneumonia	7	15	3	4	28-60	40
Falls	5	11	2	3	32-46	41
Sepsis	3	6	3	0	46-56	52
Complications of general anesthesia	2	4	0	2	12-15	14
Intestinal obstruction	1	2	1	0	28	28
Starvation	1	2	1	0	15	15
Drowning	1	2	0	1	3	3
Accident	1	2	1	0	25	25
Suicide	1	2	1	0	30	30
Total	48	100	24	24	3-60	40

84% of the population of the United States in 1980 survived to the age of sixty years. The median survival for the 1980 U.S. population is estimated to be seventy-eight years, some twenty-two years greater than the estimate for the cohort of patients with fibrodysplasia ossificans progressiva.

The most common causes of death in patients with fibrodysplasia ossificans progressiva were cardiorespiratory failure from thoracic insufficiency syndrome (54%; twenty-six [twelve male and fourteen female] individuals, with a median age of forty-two years; range, eight to fifty-eight years), pneumonia (15%; seven [three male and four female] individuals, with a median age of forty years; range, twenty-eight to sixty years), and complications of falls (11%; five [two male and three female] individuals with a median age of forty-one years; range, thirty-two to forty-six years) (Table I). All deaths from falls were ascribed to complications of head injuries, a well-known morbidity factor in patients with fibrodysplasia ossificans progressiva³. Surprisingly, there were no reports of complete autopsies in any patient.

The median age at the time of death was thirty-nine years for those with no clearly established cause of death, forty-two years for those who had cardiorespiratory failure from thoracic insufficiency syndrome, forty years for those who had pneumonia, and forty-one years for those who died from a fall. Deaths from cardiorespiratory failure and pneumonia occurred within a fifty-year range and a thirty-two-year range, respectively, while deaths from falls occurred in a narrower fourteen-year age range (between thirty-two and forty-six years; $p = 0.008$) (Table I), most likely reflecting an age range in which some patients were still able to walk or able to transfer with assistance but were at greatest risk of falls from progressive instability.

Discussion

There are two major findings of our study: (1) fibrodysplasia ossificans progressiva is not only an extremely disabling disease but also a condition of considerably shortened lifespan, and (2) the most common cause of death in patients with fibrodysplasia ossificans progressiva is cardiorespiratory

failure from thoracic insufficiency syndrome. Numerous factors contribute to the development of thoracic insufficiency syndrome in patients with fibrodysplasia ossificans progressiva, including costovertebral malformations with early orthotopic ankylosis of the costovertebral joints; fusion of ribs; ossification of intercostal muscles, paravertebral muscles, and aponeuroses; and progressive spinal deformity including kyphoscoliosis or thoracic lordosis (Fig. 3)¹¹. Scoliosis surgery is contraindicated in most patients as operative intervention exacerbates the disease process¹².

Thoracic insufficiency syndrome and severe restrictive lung disease occur in all individuals with fibrodysplasia ossificans progressiva¹¹. In a large prospective study of young, active patients with fibrodysplasia ossificans progressiva, echocardiograms were still normal, but 40% of the patients had electrocardiographic evidence of right ventricular dysfunction, including right ventricular hypertrophy, P pul monale, incomplete right bundle branch block, and right ventricular strain pattern¹³. These changes are indicative of cor pulmonale.

In our study, individuals who died of cardiorespiratory failure had a terminal course that was similar to patients with severe pulmonary hypertension. These patients typically present with progressive dyspnea, edema, elevated jugular venous pressure with marked V waves, right ventricular heave, hypoxemia, and electrocardiographic evidence of right ventricular dysfunction¹⁴. Pulmonary hypertension, cor pulmonale, and cardiorespiratory failure are all terminal features of prolonged alveolar hypoventilation due to the severe thoracic insufficiency syndrome¹⁴. In one postmortem report, gross right ventricular hypertrophy was confirmed in a patient who died of respiratory failure after presenting with all of the classic signs and symptoms of severe pulmonary hypertension and electrocardiographic evidence of right ventricular dysfunction¹⁵. These studies indicate that periodic electrocardiographic evaluations of individuals with fibrodysplasia ossificans progressiva are useful to monitor the development of right ventricular dysfunction^{16,17}. The patients with electrocardiographic evidence of right ventricular dys-




Fig. 3
Working model of the pathophysiology of thoracic insufficiency syndrome and cardiorespiratory failure in patients with fibrodysplasia ossificans progressiva.

function require additional care to guard against catastrophic hemodynamic collapse that can occur in the setting of severe pulmonary hypertension.

Nearly all individuals who died of cardiorespiratory failure from thoracic insufficiency syndrome had used supplemental oxygen at home in the week prior to their death despite warnings to the contrary. In several individuals who were hospitalized prior to their death, carbon dioxide levels were elevated in the arterial blood. Patients with end-stage thoracic insufficiency syndrome are chronically hypercarbic and have lost an important component of respiratory drive associated with acute hypercarbia^{11,18}. In such a setting, the therapeutic use of supplemental oxygen can lead to the further loss of respiratory drive and a subsequent rapid demise¹¹. Thus, patients with fibrodysplasia ossificans progressiva should not use unmonitored oxygen therapy, especially during sleep. However, individuals who have thoracic insufficiency may benefit from noninvasive mechanical methods to increase oxygen saturation and relieve chronic hypercarbia¹¹.

While pulmonary hypertension in patients with fibrodysplasia ossificans progressiva has been largely attributed to chronic hypoventilation from thoracic insufficiency, there is an intriguing possibility that the dysregulation of the BMP pathway that causes fibrodysplasia ossificans progressiva¹⁹⁻²³ may play a direct role in the pathogenesis of pulmonary hypertension. While the mutation in ACVR1/ALK2 in patients

with fibrodysplasia ossificans progressiva leads to activation of BMP signaling^{4,20,24}, the BMPR2 mutations in primary pulmonary hypertension are generally thought to lead to downregulation of BMP signaling²⁵⁻³⁴. However, given the functional redundancy among BMP type-II receptors, it is increasingly recognized that loss of Smad signaling^{35,36}. In fact, evidence suggests that overactive BMP signaling, as in fibrodysplasia ossificans progressiva, may play a central role in the development of hypoxia-induced pulmonary hypertension³⁷. In addition, recent studies have shown that activating mutations of ACVR1, similar to those seen in patients with fibrodysplasia ossificans progressiva, lead to paradoxical changes in the expression of BMP type-II receptors and in downstream BMP signaling in affected cells³⁸. Thus, whether cardiorespiratory mortality in patients with fibrodysplasia ossificans progressiva is due to restrictive chest-wall disease alone or has a pulmonary vascular component resulting directly from aberrant ACVR1/ALK2 signaling is an intriguing area worthy of further investigation. Genetic knock-in animal models of fibrodysplasia ossificans progressiva are presently under development and may help to clarify this important question. Noninvasive echocardiographic studies late in the course of the disease coupled with thorough histopathologic evaluation of the myocardium and lung parenchyma following death in patients with fibrodysplasia ossificans progressiva may also help to resolve this critical issue.


Despite the decreased median lifespan of these patients, there is wide variation in lifespan, even among individuals with the identical ACVR1/ALK2 receptor mutation. This observation suggests that other genes in the BMP signaling pathway or other pathways might modify the enhanced signaling output of the mutant receptor and be responsible in part for lifespan variation. Additionally, environmental factors also play an important role in the postnatal phenotype of fibrodysplasia ossificans progressiva. In a recent study of three pairs of monozygotic twins with fibrodysplasia ossificans progressiva, each pair of twins had identical congenital toe malformations³⁹. However, postnatal heterotopic ossification and chest wall deformity varied greatly between the twins in each pair, depending on life history and environmental exposure. Thus, the classic fibrodysplasia ossificans progressiva mutation strongly influences the phenotype of the condition during prenatal development, while environmental factors strongly influence postnatal progression of the disease³⁹.

In a recent study, we observed genotype-phenotype correlations between some exceedingly rare ACVR1/ALK2 mutations and the age of onset of heterotopic ossification or in embryonic patterns of skeletal development⁴⁰. However, <2% of individuals with fibrodysplasia ossificans progressiva have phenotypic and genotypic variants⁴⁰. Importantly, all of the individuals in our study had classic fibrodysplasia ossificans progressiva. In those for whom banked DNA was available for testing, the R206H mutation in ACVR1/ALK2 was confirmed. Thus, phenotype-genotype correlations were not possible.

Greater awareness of fibrodysplasia ossificans progressiva among clinicians, earlier diagnosis of the condition, and educational programs to avoid iatrogenic harm strongly suggest that mortality data should be stable, and perhaps even improving^{3,41-43}. Importantly, the recent discovery of the fibrodysplasia ossificans progressiva gene has provided critical insight into the development of medications that have the potential to alter the natural history of this disorder^{38,44,45}.

In summary, there are two major findings of our study: (1) fibrodysplasia ossificans progressiva is not only an extremely disabling disease but also a condition of considerably shortened lifespan, and (2) the most common cause of death in patients with fibrodysplasia ossificans progressiva is cardiorespiratory failure from thoracic insufficiency syndrome. This study has important implications for understanding the natural history of the disease, for establishing baseline mortality profiles for life-care planning, and for designing studies to better understand the complex pathophysiology of cardiorespiratory failure in fibrodysplasia ossificans progressiva.

Appendix

 A table listing all patients and their cause of death is available with the electronic version of this article on our web site at jbjs.org (go to the article citation and click on "Supporting Data"). ■

Frederick S. Kaplan, MD
Department of Orthopaedic Surgery,
Hospital of the University of Pennsylvania,

Silverstein-2, 3400 Spruce Street,
Philadelphia, PA 19104.
E-mail address for F.S. Kaplan:
Frederick.Kaplan@uphs.upenn.edu

Michael A. Zasloff, MD, PhD
Department of Translational Sciences,
Georgetown University Medical Center,
Washington, DC 20057

Joseph A. Kitterman, MD
Department of Pediatrics and
Cardiovascular Research Institute, U-503; Box-0734,
the University of California at San Francisco,
San Francisco, CA 94143-0734

Eileen M. Shore, PhD
Department of Orthopaedic Surgery,
University of Pennsylvania School of Medicine,
424 Stemmler Hall,
36th and Hamilton Walk,
Philadelphia, PA 19104

Charles C. Hong, MD, PhD
Departments of Medicine and Pharmacology
and Center for Inherited Heart Disease,
Vanderbilt University School of Medicine,
2220 Pierce Avenue,
383 Preston Research Building,
Nashville, TN 37232

David M. Rocke, PhD
Department of Public Health Sciences,
University of California at Davis,
One Shields Avenue,
Davis, CA 95616

References

- Kaplan FS, Glaser DL, Shore EM, Deirmengian GK, Gupta R, Delai P, Morhart R, Smith R, Le Merrer M, Rogers JG, Connor M, Kitterman JA. The phenotype of fibrodysplasia ossificans progressiva. *Clin Rev Bone Miner Metab*. 2005;3:4:183-8.
- Kaplan FS, Groppe J, Pignolo RJ, Shore EM. Morphogen receptor genes and metamorphogenesis: skeleton keys to metamorphosis. *Ann N Y Acad Sci*. 2007; 1116:113-33.
- Kaplan FS, LeMerrer M, Glaser DL, Pignolo RJ, Goldsby RE, Kitterman JA, Groppe J, Shore EM. Fibrodysplasia ossificans progressiva. *Best Pract Res Clin Rheumatol*. 2008;22:191-205.
- Shore EM, Xu M, Feldman GJ, Fenstermacher DA, Cho T-J, Choi IH, Connor JM, Delai P, Glaser DL, Le Merrer M, Morhart R, Rogers JG, Smith R, Triffitt JT, Urtizberea JA, Zasloff M, Brown MA, Kaplan FS. A recurrent mutation in the BMP type I receptor ACVR1 causes inherited and sporadic fibrodysplasia ossificans progressiva. *Nat Genet*. 2006;38:525-7.
- Shore EM, Feldman GJ, Xu M, Kaplan FS. The genetics of fibrodysplasia ossificans progressiva. *Clin Rev Bone Miner Metab*. 2005;3:4:201-4.
- Cohen RB, Hahn GV, Tabas J, Peeper J, Levitz CL, Sando A, Sando N, Zasloff M, Kaplan FS. The natural history of heterotopic ossification in patients who have fibrodysplasia ossificans progressiva. A study of forty-four patients. *J Bone Joint Surg Am*. 1993;75:215-9.
- Rocke DM, Zasloff M, Peeper J, Cohen RB, Kaplan FS. Age- and joint-specific risk of initial heterotopic ossification in patients who have fibrodysplasia ossificans progressiva. *Clin Orthop Relat Res*. 1994;301:243-8.
- Glaser DL, Kaplan FS. Treatment considerations for the management of fibrodysplasia ossificans progressiva. *Clin Rev Bone Miner Metab*. 2005;3:4:243-50.
- R Development Core Team. R: A language and environment for statistical computing. Version 2.8.1. Vienna: R Foundation for Statistical Computing. <http://www.r-project.org>. Accessed 2009 Jan 8.
- Therneau T, Lumley T. Survival: survival analysis including penalized likelihood. R package version 2.34-1. <http://CRAN.R-project.org/package=survival>. Accessed 2009 Jan 8.
- Kaplan FS, Glaser DL. Thoracic insufficiency syndrome in patients with fibrodysplasia ossificans progressiva. *Clin Rev Bone Miner Metab*. 2005;3:4:213-6.
- Shah PB, Zasloff MA, Drummond D, Kaplan FS. Spinal deformity in patients who have fibrodysplasia ossificans progressiva. *J Bone Joint Surg Am*. 1994;76: 1442-50.
- Kussmaul WG, Esmail AN, Sagar Y, Ross J, Gregory S, Kaplan FS. Pulmonary and cardiac function in advanced fibrodysplasia ossificans progressiva. *Clin Orthop Relat Res*. 1998;346:104-9.
- D'Alonzo GE, Barst RJ, Ayres SM, Bergofsky EH, Brundage BH, Detre KM, Fishman AP, Goldring RM, Groves BM, Kernis JT, Levy PS, Pietra GG, Reid LM, Reeves JT, Rich S, Veim CE, Williams GW, Wu M. Survival in patients with primary pulmonary hypertension. Results from a national prospective registry. *Ann Intern Med*. 1991;115:343-9.
- Orme S, Underwood JC, McCloskey E, Hendra TJ. An unusual cause of pulmonary hypertension and right heart failure. *Postgrad Med J*. 1998;74:697-8.
- Buhain WJ, Rammohan G, Berger HW. Pulmonary function in myositis ossificans progressiva. *Am Rev Respir Dis*. 1974;110:333-7.
- Connor JM, Evans CC, Evans DAP. Cardiopulmonary function in fibrodysplasia ossificans progressiva. *Thorax*. 1981;36:419-23.

18. Campbell RM, Smith MD. Thoracic insufficiency syndrome and exotic scoliosis. *J Bone Joint Surg Am.* 2007;89 Suppl 1:108-22.
19. Ahn J, Serrano de la Peña L, Shore EM, Kaplan FS. Paresis of a bone morphogenetic protein-antagonist response in a genetic disorder of heterotopic skeletogenesis. *J Bone Joint Surg Am.* 2003;85:667-74.
20. Billings PC, Fiori JL, Bentwood JL, O'Connell MP, Jiao X, Nussbaum B, Caron RJ, Shore EM, Kaplan FS. Dysregulated BMP signaling and enhanced osteogenic differentiation of connective tissue progenitor cells from patients with fibrodysplasia ossificans progressiva (FOP). *J Bone Miner Res.* 2008;23:305-13.
21. Fiori JL, Billings PC, de la Peña LS, Kaplan FS, Shore EM. Dysregulation of the BMP-p38 MAPK signaling pathway in cells from patients with fibrodysplasia ossificans progressiva (FOP). *J Bone Miner Res.* 2006;21:902-9.
22. de la Peña LS, Billings PC, Fiori JL, Ahn J, Kaplan FS, Shore EM. Fibrodysplasia ossificans progressiva (FOP), a disorder of ectopic osteogenesis, misregulates cell surface expression and trafficking of BMPRIA. *J Bone Miner Res.* 2005;20:1168-76.
23. Shafritz AB, Shore EM, Gannon FH, Zasloff MA, Taub R, Muenke M, Kaplan FS. Overexpression of an osteogenic morphogen in fibrodysplasia ossificans progressiva. *N Engl J Med.* 1996;335:555-61.
24. Shen Q, Little SC, Xu M, Haupt J, Ast C, Katagiri T, Mundlos S, Seemann P, Kaplan FS, Mullins MC, Shore EM. The fibrodysplasia ossificans progressiva R206H ACVR1 mutation activates BMP-independent chondrogenesis and zebrafish embryo ventralization. *J Clin Invest.* 2009;119:3462-72.
25. Atkinson C, Stewart S, Upton PD, Machado R, Thomson JR, Trembath RC, Morrell NW. Primary pulmonary hypertension is associated with reduced pulmonary vascular expression for type II bone morphogenetic protein receptor. *Circulation.* 2002;105:1672-8.
26. Du L, Sullivan CC, Chu D, Cho AJ, Kido M, Wolf PL, Yuan J, Deutsch R, Jamieson SW, Thistlethwaite PA. Signaling molecules in nonfamilial pulmonary hypertension. *N Engl J Med.* 2003;348:500-9.
27. Eblaghie MC, Reedy M, Oliver T, Mishina Y, Hogan BL. Evidence that autocrine signaling through Bmpr1a regulates the proliferation, survival and morphogenetic behavior of distal lung epithelial cells. *Dev Biol.* 2006;291:67-82.
28. Machado RD, Aldred MA, James V, Harrison RE, Patel B, Schwalbe EC, Gruenig E, Janssen B, Koehler R, Seeger W, Eickelberg O, Olschewski H, Elliott CG, Glissmeyer E, Carlquist J, Kim M, Torbicki A, Fijalkowska A, Szweczyk G, Parma J, Abramowicz MJ, Galie N, Morisaki H, Kyotani S, Nakanishi N, Morisaki T, Humbert M, Simonneau G, Sitbon O, Soubrier F, Coulet F, Morrell NW, Trembath RC. Mutations of the TGF-beta type II receptor BMPR2 in pulmonary arterial hypertension. *Hum Mutat.* 2006;27:121-32.
29. Morrell NW. Pulmonary hypertension due to BMPR2 mutation: a new paradigm for tissue remodeling? *Proc Am Thoracic Soc.* 2006;3:680-6.
30. Rabinovitch M. Molecular pathogenesis of pulmonary arterial hypertension. *J Clin Invest.* 2008;118:2372-9.
31. Southwood M, Jeffery TK, Yang X, Upton PD, Hall SM, Atkinson C, Haworth SG, Stewart S, Reynolds PN, Long L, Trembath RC, Morrell NW. Regulation of bone morphogenetic protein signalling in human pulmonary vascular development. *J Pathol.* 2008;214:85-95.
32. International PPH Consortium, Lane KB, Machado RD, Pauciulo MW, Thomson JR, Phillips JA 3rd, Loyd JE, Nichols WC, Trembath RC. Heterozygous germline mutations in BMPR2, encoding a TGF-beta receptor, cause familial primary pulmonary hypertension. *Nat Genet.* 2000;26:81-4.
33. Thomson JR, Machado RD, Pauciulo MW, Morgan NV, Humbert M, Elliott GC, Ward K, Yacoub M, Mikhail G, Rogers P, Newman J, Wheeler L, Higenbottam T, Gibbs JS, Egan J, Crozier A, Peacock A, Allcock R, Corris P, Loyd JE, Trembath RC, Nichols WC. Sporadic primary pulmonary hypertension is associated with germline mutations of the gene encoding BMPR-II, a receptor member of the TGF-beta family. *J Med Genet.* 2000;37:741-5.
34. Yang X, Long L, Southwood M, Rudarakanchana N, Upton PD, Jeffery TK, Atkinson C, Chen H, Trembath RC, Morrell NW. Dysfunctional Smad signaling contributes to abnormal smooth muscle cell proliferation in familial pulmonary arterial hypertension. *Circ Res.* 2005;96:1053-63.
35. Yu PB, Beppu H, Kawai N, Li E, Bloch KD. Bone morphogenetic protein (BMP) type II receptor deletion reveals BMP ligand-specific gain of signaling in pulmonary artery smooth muscle cells. *J Biol Chem.* 2005;280:24443-50.
36. Yu PB, Deng DY, Beppu H, Hong CC, Lai C, Hoeng SA, Kawai N, Bloch KD. Bone morphogenetic protein (BMP) type II receptor is required for BMP-mediated growth arrest and differentiation in pulmonary artery smooth muscle cells. *J Biol Chem.* 2008;283:3877-88.
37. Frank DB, Abtahi A, Yamaguchi DJ, Manning S, Shyr Y, Pozzi A, Baldwin HS, Johnson JE, de Caestecker MP. Bone morphogenetic protein 4 promotes pulmonary vascular remodeling in hypoxic pulmonary hypertension. *Circ Res.* 2005;97:496-504.
38. Yu P, Deng DY, Lai CS, Hong CC, Cuny GD, Bouxsein ML, Hong DW, McManus PM, Katagiri T, Sachidanandan C, Kamiya N, Fukuda T, Mishina Y, Peterson RT, Bloch KD. BMP type I receptor inhibition reduces heterotopic [corrected] ossification. *Nat Med.* 2008;14:1363-9. Erratum in: *Nat Med.* 2009;15:117.
39. Hebel N, Shore EM, Kaplan FS. Three pairs of monozygotic twins with fibrodysplasia ossificans progressiva: the role of environment in the progression of heterotopic ossification. *Clin Rev Bone Miner Metab.* 2005;3:4:205-8.
40. Kaplan FS, Xu M, Seemann P, Connor JM, Glaser DL, Carroll L, Delai P, Fastnacht-Urban E, Forman SJ, Gillesen-Kaesbach G, Hoover Fong J, Köster B, Pauli RM, Reardon W, Zaidi S-A, Zasloff M, Morhart R, Mundlos S, Groppe J, Shore EM. Classic and atypical fibrodysplasia ossificans progressiva (FOP) phenotypes are caused by mutations in the bone morphogenetic protein (BMP) type I receptor ACVR1. *Hum Mutat.* 2009;30:379-90.
41. Hair MS, Peeper JL. The international fibrodysplasia ossificans progressiva association. *Clin Rev Bone Miner Metab.* 2005;3:4:267-9.
42. Kaplan FS, Xu M, Glaser DL, Collins F, Connor M, Kitterman J, Sillence D, Zackai E, Ravitsky V, Zasloff M, Ganguly A, Shore EM. Early diagnosis of fibrodysplasia ossificans progressiva. *Pediatrics.* 2008;121:e1295-300.
43. Kitterman JA, Kantanie S, Rocke DM, Kaplan FS. Iatrogenic harm caused by diagnostic errors in fibrodysplasia ossificans progressiva. *Pediatrics.* 2005;116:e654-61.
44. Kaplan FS. The key to the closet is the key to the kingdom: a common lesson of rare diseases. *Orphan Disease Update.* 2006;24:1-9.
45. Kaplan FS, Groppe J, Shore EM. When one skeleton is enough: approaches and strategies for the treatment of fibrodysplasia ossificans progressiva (FOP). *Drug Discovery Today: Therapeutic Strategies.* 2008;5:255-62.