

Homework 4 Comments

David M. Roche

February 11, 2025

- For document in Word or the equivalent, things will look funny unless you change to a monospace font. Default fonts have characters of variable width and different spaces between characters. These are called proportionally spaced. The standard monospace font in Windows is Courier New. For Macs, the traditional monospace font is SF Mono Regular, though there are others. If you copy from R to a document with a monospaced font, it does not change the appearance.
- Choose No Spacing on the Home tab.
- Highlight the whole document (at the start or later) with Ctrl-a.
- Choose Courier New and 11 or 12 points. There are 72 points to the inch.
- Choose paragraph on the home tab. Make sure line spacing is single and spacing before/after is 0

- If a variable is coded with numbers, such as 1, 2, 3, . . . , then ask yourself if it is a measurement of something. If not, then make it a factor.
- In a clinical study, weight is a numeric variable, systolic blood pressure is a numerical variable, but patient number is not a numerical variable and needs to be a factor.

- If you have a document with code and output, and if the code would not produce the output you included, what am I to think?
- Do not type in numbers which are already in a file you have loaded or should have loaded. This leads to frequent errors.

- You should almost never use the equal variance t-test.
- If the variances of two groups really are equal, the performance of the equal variance t-test is a little better than that of the Welch t-test which does not make that assumption.
- If the variances truly are not equal, the performance of the equal variance t-test will be worse, sometime catastrophically worse.
- It is often hard to tell if the variances really are equal
- This makes the equal variance test a bad gamble, with small possible gain, and large possible loss.
- For more than two groups, there is no good alternative to the equal variance analysis, but check if there are large variations in the variance.

- Statistical tests for normality, equality of variance, serial correlation and others are often not very informative.
- They generally have low statistical power, which is the chance of a significant result if the equality etc. is not true.
- Conditional use of these tests is often bad: test for equal variance; if $p < 0.05$ use the Welch t-test; otherwise use the equal variance t-test. Usually, these compound procedures perform worse than the Welch test without a pre-test.
- An even worse procedure is testing for normality: if significant, use a rank test; if not, use a t-test.
- Procedures like the Welch t-test or linear regression often work well even if the exact assumptions fail. Exact assumptions are always false! Approximately true is usually good enough.

If there is an outlier from the plots, as there is in 4.5/4.2 in the homework, it would make sense to see what the analysis looks like without that point. Similarly, in 4.3/3.15 the significance of the numeric order of the run is disturbing, even with randomization.

Don't edit the output too much: `summary(lm)` is much better than `print(lm)`.

```
> print(lm(y~method+run, data = yield))
```

Call:

```
lm(formula = y ~ method + run, data = yield)
```

Coefficients:

(Intercept)	methodB	run
43.9459	20.0789	0.7723


```
> summary(lm(y~method+run, data = yield))
```

Call:

```
lm(formula = y ~ method + run, data = yield)
```

Residuals:

Min	1Q	Median	3Q	Max
-9.9589	-2.9716	-0.3824	4.1236	8.9689

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)	
(Intercept)	43.9459	2.7972	15.711	1.48e-11	***
methodB	20.0789	2.3984	8.372	1.96e-07	***
run	0.7723	0.2080	3.714	0.00173	**

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 5.36 on 17 degrees of freedom

Multiple R-squared: 0.828, Adjusted R-squared: 0.8077

F-statistic: 40.91 on 2 and 17 DF, p-value: 3.181e-07